



THE UNIVERSITY of EDINBURGH

Edinburgh Research Explorer

ARIA pharmacy 2018 "Allergic rhinitis care pathways for community pharmacy"

Citation for published version:

Bosnic-Anticevich, S, Costa, E, Menditto, E, Lourenço, O, Novellino, E, Bialek, S, Briedis, V, Buonaiuto, R, Chrystyn, H, Cvetkovski, B, Capua, SD, Kritikos, V, Mair, A, Orlando, V, Paulino, E, Salimäki, J, Söderlund, R, Tan, R, Williams, DM, Wroczynski, P, Agache, I, Ansotegui, IJ, Anto, JM, Bedbrook, A, Bachert, C, Bewick, M, Bindeslev-Jensen, C, Brozek, J, Canonica, GW, Cardona, V, Carr, W, Casale, T, Chavannes, NH, Correia de Sousa, J, Cruz, AA, Czarlewski, W, De Carlo, G, Demoly, P, Devillier, P, Dykewicz, MS, Gaga, M, El-Gamal, Y, Fonseca, J, Fokkens, WJ, Guzmán, MA, Haahtela, T, Hellings, PW, Illario, M, Ivancevich, JC, Just, J, Kaidashev, I, Khaitov, M, Khaltaev, N, Keil, T, Klimek, L, Kowalski, ML, Kuna, P, Kvedariene, V, Larenas-Linnemann, D, Laune, D, Le, LTT, Carlsen, KCL, Mahboub, B, Maier, D, Malva, J, Manning, P, Morais-Almeida, M, Mösges, R, Mullol, J, Münter, L, Murray, R, Naclerio, R, Namazova-Baranova, L, Nekam, K, Nyembue, TD, Okubo, K, O'Hehir, RE, Ohta, K, Okamoto, Y, Onorato, GL, Palkonen, S, Panzner, P, Papadopoulos, NG, Park, HS, Pawankar, R, Pfaar, O, Phillips, J, Plavec, D, Popov, TA, Potter, P, Prokopakis, EP, Roller-Wirnsberger, RE, Rottem, M, Ryan, D, Samolinski, B, Sanchez-Borges, M, Schunemann, HJ, Sheikh, A, Sisul, JC, Somekh, D, Stellato, C, To, T, Todo-Bom, A, Tomazic, PV, Toppila-Salmi, S, Valero, A, Valiulis, A, Valovirta, E, Ventura, MT, Wagenmann, M, Wallace, D, Wasserman, S, Wickman, M, Yiallourous, PK, Yorgancioglu, A, Yusuf, OM, Zar, HJ, Zernotti, ME, Zhang, L, Zidarn, M, Zuberbier, T & Bousquet, J 2018, 'ARIA pharmacy 2018 "Allergic rhinitis care pathways for community pharmacy"', *Allergy*. <https://doi.org/10.1111/all.13701>

Digital Object Identifier (DOI):

[10.1111/all.13701](https://doi.org/10.1111/all.13701)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Allergy

Publisher Rights Statement:

This is the author's peer-reviewed manuscript as accepted for publication.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Download date: 08. May. 2023



PROF. OLGA LOURENÇO (Orcid ID : 0000-0002-8401-5976)

MS. RACHEL TAN (Orcid ID : 0000-0001-5171-4036)

DR. IOANA AGACHE (Orcid ID : 0000-0001-7994-364X)

PROF. CLAUS BACHERT (Orcid ID : 0000-0003-4742-1665)

DR. VICTORIA CARDONA (Orcid ID : 0000-0003-2197-9767)

PROF. THOMAS B CASALE (Orcid ID : 0000-0002-3149-7377)

PROF. JAIME CORREIA DE SOUSA (Orcid ID : 0000-0001-6459-7908)

PROF. PHILIPPE DEVILLIER (Orcid ID : 0000-0003-4107-8317)

PROF. MAREK LESZEK KOWALSKI (Orcid ID : 0000-0002-8442-2774)

DR. DÉsirÉE ERLINDA LARENAS-LINNE MANN (Orcid ID : 0000-0002-5713-5331)

PROF. KARIN C. LØDRUP CARLSEN (Orcid ID : 0000-0002-9257-1198)

PROF. RALPH MOSGES (Orcid ID : 0000-0002-1928-810X)

DR. JOAQUIM MULLOL (Orcid ID : 0000-0003-3463-5007)

DR. KEN OHTA (Orcid ID : 0000-0001-9734-4579)

DR. NIKOLAOS G PAPADOPOULOS (Orcid ID : 0000-0002-4448-3468)

PROF. HAE-SIM PARK (Orcid ID : 0000-0003-2614-0303)

PROF. OLIVER PFAAR (Orcid ID : 0000-0003-4374-9639)

DR. DERMOT RYAN (Orcid ID : 0000-0002-4115-7376)

MS. FABIENNE PORTEJOIE (Orcid ID : 0000-0001-9226-7762)

Article type : Position Paper

Editor : Luo Zhang

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/all.13701

This article is protected by copyright. All rights reserved.

ARIA pharmacy 2018

“Allergic rhinitis care pathways for community pharmacy”

AIRWAYS ICPs initiative (European Innovation Partnership on Active and Healthy Ageing, DG CNECT and DG Santé)

POLLAR (Impact of Air Pollution on asthma and rhinitis)

GARD Demonstration project

S Bosnic-Anticevich (1), E Costa (2), E Menditto (3), O Lourenço (4), E Novellino (5), S Bialek (6), V Briedis (7), R Buonaiuto (8), H Chrystyn (9), B Cvetkovski (1), S Di Capua (10), V Kritikos (1), A Mair (11), V Orlando (3), E Paulino (12), J Salimäki (13), R Söderlund (14), R Tan (1), DM Williams (15), P Wroczynski (6), I Agache (16), IJ Ansotegui (17), JM Anto (18-21), A Bedbrook (22), C Bachert (23), M Bewick (24), C Bindselev-Jensen (25), J Brozek (26), GW Canonica (27), V Cardona (28), W Carr (29), T Casale (30), NH Chavannes (31), J Correia de Sousa (32), AA Cruz (33), W Czarlewski (34), G De Carlo (35), P Demoly (36, 37), P Devillier (38), MS Dykewicz (39), M Gaga (40), Y El-Gamal (41), J Fonseca (42), WJ Fokkens (43), MA Guzmán (44), T Haahtela (45), PW Hellings (46), M Illario (47), JC Ivancevich (48), J Just (49), I Kaidashev (50), M Khaitov (51), N Khaltaev (52), T Keil (53), L Klimek (54), ML Kowalski (55), P Kuna (56), V Kvedariene (57), D Larenas-Linnemann (58), D Laune (59), LTT Le (60), KC Lodrup Carlsen (61), B Mahboub (62), D Maier (63), J Malva (64), P Manning (65), M Morais-Almeida (66), R Mösges (67), J Mullol (68), L Münter (69), R Murray (70), R Naclerio (71), L Namazova-Baranova (72), K Nekam (73), TD Nyembue (74), K Okubo (75), RE O’Hehir (76), K Ohta (77), Y Okamoto (78), GL Onorato (22), S Palkonen (35), P Panzner (79), NG Papadopoulos (80), HS Park (81), R Pawankar (82), O Pfaar (83), J Phillips (84), D Plavec (85), TA Popov (86), P Potter (87), EP Prokopakis (88), RE Roller-Wirnsberger (89), M Rottem (90), D Ryan (91), B Samolinski (92), M Sanchez-Borges (93), HJ Schunemann (26), A Sheikh (94), JC Sisul (95), D Somekh (96), C Stellato (97), T To (98), A Todo-Bom (99), PV Tomazic (100), S Toppila-Salmi (45), A Valero (101), A Valiulis (102), E Valovirta (103), MT Ventura (104), M Wagenmann (105), D Wallace (106), S Wasserman (107), M Wickman (108), PK Yiallourous (109), A Yorgancioglu (110), OM Yusuf (111), HJ Zar (112), ME Zernotti (113), L Zhang (114), M Zidarn (115), T Zuberbier (116), J Bousquet J (22, 116, 117, 118)

1. Woolcock Institute of Medical Research, University of Sydney and Sydney Local Health District, Glebe, NSW, Australia.
2. UCIBIO, REQUINTE, Faculty of Pharmacy and Competence Center on Active and Healthy Ageing of University of Porto (AgeUPNetWork), University of Porto, Portugal
3. CIRFF, Federico II University, Naples, Italy.

- Accepted Article
4. Faculty of Health Sciences and CICS – UBI, Health Sciences Research Centre, University of Beira Interior, Covilhã, Portugal.
 5. Director of Department of Pharmacy of University of Naples Federico II, Naples, Italy.
 6. Department of Biochemistry and Clinical Chemistry, Faculty of Pharmacy with the Division of Laboratory Medicine, Warsaw Medical University, Warsaw, Poland
 7. Head of Department of Clinical Pharmacy of Lithuanian University of Health Sciences, Kaunas, Lithuania.
 8. Pharmacist, Municipality Pharmacy, Sarno, Italy.
 9. RiRL, 5a Coles Lane, Oakington, Cambridge, UK.
 10. Farmacie Dei Golfi Group, Massa Lubrense, Italy.
 11. DG for Health and Social Care, Scottish Government, Edinburgh, UK.
 12. Farmacias Holon, Lisbon, Portugal.
 13. Association of Finnish Pharmacies.
 14. Department of Nephrology and Endocrinology, Karolinska University Hospital, Stockholm, Sweden.
 15. Eshelman School of Pharmacy, University of North Carolina, Chapel Hill, NC, USA.
 16. Transylvania University Brasov, Brasov, Romania.
 17. Department of Allergy and Immunology, Hospital Quirón Bizkaia, Erandio, Spain.
 18. ISGlobAL, Centre for Research in Environmental Epidemiology (CREAL), Barcelona, Spain.
 19. IMIM (Hospital del Mar Research Institute), Barcelona, Spain.
 20. CIBER Epidemiología y Salud Pública (CIBERESP), Barcelona, Spain.
 21. Universitat Pompeu Fabra (UPF), Barcelona, Spain.
 22. MACVIA-France, Fondation partenariale FMC VIA-LR, Montpellier, France.
 23. BUpper Airways Research Laboratory, ENT Dept, Ghent University Hospital, Ghent, Belgium.
 24. iQ4U Consultants Ltd, London, UK.
 25. Department of Dermatology and Allergy Centre, Odense University Hospital, Odense Research Center for Anaphylaxis (ORCA), Odense, Denmark.
 26. Department of Health Research Methods, Evidence, and Impact, Division of Immunology and Allergy, Department of Medicine, McMaster University, Hamilton, ON, Canada.
 27. Personalized Medicine Clinic Asthma & Allergy, Humanitas University, Humanitas Research Hospital, Rozzano, Milan, Italy
 28. Allergy Section, Department of Internal Medicine, Hospital Vall 'dHebron & ARADyAL research network, Barcelona, Spain.
 29. Allergy and Asthma Associates of Southern California, Mission Viejo, CA, USA.
 30. Division of Allergy/Immunology, University of South Florida, Tampa, Fla.
 31. Department of Public Health and Primary Care, Leiden University Medical Center, Leiden, The Netherlands

This article is protected by copyright. All rights reserved.

- Accepted Article
32. Life and Health Sciences Research Institute (ICVS), School of Medicine, University of Minho, Braga, Portugal; ICVS/3B's, PT Government Associate Laboratory, Braga/Guimarães, Portugal.
 33. ProAR – Nucleo de Excelencia em Asma, Federal University of Bahia, Brasil and WHO GARD Planning Group, Brazil.
 34. Medical Consulting Czarlewski, Levallois, France.
 35. EFA European Federation of Allergy and Airways Diseases Patients' Associations, Brussels, Belgium
 36. Department of Respiratory Diseases, Montpellier University Hospital, France.
 37. Epidemiology of Allergic and Respiratory Diseases, Department Institute Pierre Louis of Epidemiology and Public Health, INSERM and UPMC Sorbonne Université, Medical School Saint Antoine, Paris, France
 38. Laboratoire de Pharmacologie Respiratoire UPRES EA220, Hôpital Foch, Suresnes Université Versailles Saint-Quentin, Université Paris Saclay, France.
 39. Section of Allergy and Immunology, Saint Louis University School of Medicine, Saint Louis, Missouri, USA.
 40. ERS President 2017-2018, Athens Chest Hospital, 7th Resp Med Dept and Asthma Center, Athens, Greece.
 41. Pediatric Allergy and Immunology Unit, Children's hospital, Ain Shams University, Cairo, Egypt.
 42. CINTESIS, Center for Research in Health Technologies and Information Systems, Faculdade de Medicina da Universidade do Porto, Porto, Portugal and MEDIDA, Lda, Porto, Portugal.
 43. Department of Otorhinolaryngology, Academic Medical Centre, Amsterdam, the Netherlands.
 44. Immunology and Allergy Division, Clinical Hospital, University of Chile, Santiago, Chile.
 45. Skin and Allergy Hospital, Helsinki University Hospital, Helsinki, and University of Helsinki, Finland.
 46. Laboratory of Clinical Immunology, Department of Microbiology and Immunology, KU Leuven, Leuven, Belgium.
 47. Division for Health Innovation, Campania Region and Federico II University Hospital Naples (R&D and DISMET) Naples, Italy.
 48. Servicio de Alergia e Immunologia, Clinica Santa Isabel, Buenos Aires, Argentina.
 49. Allergology department, Centre de l'Asthme et des Allergies Hôpital d'Enfants Armand-Trousseau (APHP); Sorbonne Universités, UPMC Univ Paris 06, UMR_S 1136, Institut Pierre Louis d'Epidémiologie et de Santé Publique, Equipe EPAR, Paris, France.
 50. Ukrainian Medical Stomatological Academy, Poltava, Ukraine.
 51. National Research Center, Institute of Immunology, Federal Medicobiological Agency, Laboratory of Molecular immunology, Moscow, Russian Federation.
 52. GARD Chairman, Geneva, Switzerland.
 53. Institute of Social Medicine, Epidemiology and Health Economics, Charité - Universitätsmedizin Berlin, Berlin, and Institute for Clinical Epidemiology and Biometry, University of Wuerzburg, Germany
 54. Center for Rhinology and Allergology, Wiesbaden, Germany.
 55. Department of Immunology and Allergy, Healthy Ageing Research Center, Medical University of Lodz, Poland.
 56. Division of Internal Medicine, Asthma and Allergy, Barlicki University Hospital, Medical University of Lodz, Poland.
 57. Faculty of Medicine, Vilnius University, Vilnius, Lithuania.

- Accepted Article
58. Center of Excellence in Asthma and Allergy, Médica Sur Clinical Foundation and Hospital, México City, Mexico.
 59. KYomed INNOV, Montpellier, France.
 60. University of Medicine and Pharmacy, Hochiminh City, Vietnam.
 61. Oslo University Hospital, Department of Paediatrics, Oslo, and University of Oslo, Faculty of Medicine, Institute of Clinical Medicine, Oslo, Norway.
 62. Department of Pulmonary Medicine, Rashid Hospital, Dubai, UAE.
 63. Biomax Informatics AG, Munich, Germany.
 64. Coimbra Institute for Clinical and Biomedical Research (iCBR), Faculty of Medicine, University of Coimbra, Portugal; Ageing@Coimbra EIP-AHA Reference Site, Coimbra, Portugal.
 65. Department of Medicine (RCSI), Bon Secours Hospital, Glasnevin, Dublin, Ireland.
 66. Allergy Center, CUF Descobertas Hospital, Lisbon, Portugal
 67. Institute of Medical Statistics, and Computational Biology, Medical Faculty, University of Cologne, Germany and CRI-Clinical Research International-Ltd, Hamburg, Germany.
 68. Rhinology Unit & Smell Clinic, ENT Department, Hospital Clínic; Clinical & Experimental Respiratory Immunoallergy, IDIBAPS, CIBERES, University of Barcelona, Spain.
 69. Danish Committee for Health Education, Copenhagen East, Denmark.
 70. Director, MedScript Ltd, Dundalk, Co Louth, Ireland.
 71. Johns Hopkins School of Medicine, Baltimore, Maryland, USA.
 72. Scientific Centre of Children's Health under the the MoH, Russia, Russian National Research Medical University named Pirogov, Moscow, Russia.
 73. Hospital of the Hospitaller Brothers in Buda, Budapest, Hungary.
 74. ENT Department, University Hospital of Kinshasa, Kinshasa, Congo.
 75. Dept of Otolaryngology, Nippon Medical School, Tokyo, Japan.
 76. Department of Allergy, Immunology and Respiratory Medicine, Alfred Hospital and Central Clinical School, Monash University, Melbourne, Victoria, Australia; Department of Immunology, Monash University, Melbourne, Victoria, Australia.
 77. National Hospital Organization Tokyo National Hospital, Tokyo, Japan.
 78. Dept of Otorhinolaryngology, Chiba University Hospital, Chiba, Japan.
 79. Department of Immunology and Allergology, Faculty of Medicine and Faculty Hospital in Pilsen, Charles University in Prague, Pilsen, Czech Republic.
 80. Division of Infection, Immunity & Respiratory Medicine, Royal Manchester Children's Hospital, University of Manchester, Manchester, UK, and Allergy Department, 2nd Pediatric Clinic, Athens General Children's Hospital "P&A Kyriakou," University of Athens, Greece.
 81. Department of Allergy and Clinical Immunology, Ajou University School of Medicine, Suwon, South Korea.
 82. Department of Pediatrics, Nippon Medical School, Tokyo, Japan.

- Accepted Article
83. Department of Otorhinolaryngology, Head and Neck Surgery, Section for Rhinology and Allergy, University Hospital Marburg, Philipps-Universität Marburg, Germany.
 84. Centre for empowering patients and communities, Faulkland, Somerset, UK.
 85. Children's Hospital Srebrnjak, Zagreb, School of Medicine, University J.J. Strossmayer, Osijek, Croatia.
 86. University Hospital 'Sv Ivan Rilski'", Sofia, Bulgaria.
 87. Allergy Diagnostic and Clinical Research Unit, University of Cape Town Lung Institute, Cape Town, South Africa.
 88. Department of Otorhinolaryngology University of Crete School of Medicine, Heraklion, Greece.
 89. Medical University of Graz, Department of Internal Medicine, Graz, Austria.
 90. Division of Allergy Asthma and Clinical Immunology, Emek Medical Center, Afula, Israel.
 91. Honorary Clinical Research Fellow, Allergy and Respiratory Research Group, The University of Edinburgh, Past President SLAAI, FACAAl, Edinburgh, UK.
 92. Department of Prevention of Environmental Hazards and Allergology, Medical University of Warsaw, Poland.
 93. Allergy and Clinical Immunology Department, Centro Médico-Docente la, Trinidad and Clínica El Avila, Altamira, Caracas, Venezuela.
 94. The Usher Institute of Population Health Sciences and Informatics, The University of Edinburgh, Edinburgh, UK.
 95. Sociedad Paraguaya de Alergia Asma e Inmunología, Paraguay.
 96. European Health Futures Forum (EHFF), Isle of Wight, UK.
 97. Department of Medicine, Surgery and Dentistry "Scuola Medica Salernitana", University of Salerno, Salerno, Italy.
 98. The Hospital for Sick Children, Dalla Lana School of Public Health, University of Toronto, Toronto, Canada.
 99. Imunoalergologia, Centro Hospitalar Universitário de Coimbra and Faculty of Medicine, University of Coimbra, Portugal.
 100. Department of ENT, Medical University of Graz, Austria.
 101. Pneumology and Allergy Department CIBERES and Clinical & Experimental Respiratory Immunology, IDIBAPS, University of Barcelona, Spain.
 102. Vilnius University Institute of Clinical Medicine, Clinic of Children's Diseases, and Institute of Health Sciences, Department of Public Health, Vilnius, Lithuania; European Academy of Paediatrics (EAP/UEMS-SP), Brussels, Belgium.
 103. Department of Lung Diseases and Clinical Immunology, University of Turku and Terveystalo allergy clinic, Turku, Finland.
 104. University of Bari Medical School, Unit of Geriatric Immunology, Bari, Italy.
 105. Dept of Otorhinolaryngology, Universitätsklinikum Düsseldorf, Germany.
 106. Nova Southeastern University, Fort Lauderdale, Florida, USA.
 107. Department of Medicine, Clinical Immunology and Allergy, McMaster University, Hamilton, Ontario, Canada.
 108. Centre for Clinical Research Sörmland, Uppsala University, Eskilstuna, Sweden.

109. Cyprus International Institute for Environmental & Public Health in Association with Harvard School of Public Health, Cyprus University of Technology, Limassol, Cyprus; Department of Pediatrics, Hospital "Archbishop Makarios III", Nicosia, Cyprus.
110. Celal Bayar University Department of Pulmonology, Manisa, Turkey.
111. The Allergy and Asthma Institute, Pakistan.
112. Department of Paediatrics and Child Health, Red Cross Children's, Hospital, and MRC Unit on Child & Adolescent Health, University of Cape Town, Cape Town, South Africa.
113. Universidad Católica de Córdoba, Córdoba, Argentina.
114. Department of Otolaryngology Head and Neck Surgery, Beijing TongRen Hospital and Beijing Institute of Otolaryngology, Beijing, China.
115. University Clinic of Respiratory and Allergic Diseases, Golnik, Slovenia.
116. Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Comprehensive Allergy Center, Department of Dermatology and Allergy, a member of GA²LEN, Berlin, Germany
117. University Hospital, Montpellier, France.
118. INSERM U 1168, VIMA : Ageing and chronic diseases Epidemiological and public health approaches, Villejuif, Université Versailles St-Quentin-en-Yvelines, UMR-S 1168, Montigny le Bretonneux, France and Euforea, Brussels, Belgium

Short title: ARIA in the pharmacy

Correspondance to

Professor Jean Bousquet

CHU Arnaud de Villeneuve, 371 Avenue du Doyen Gaston Giraud, 34295 Montpellier Cedex 5, France

Tel +33 611 42 88 47 jean.bousquet@orange.fr

Source of funding: European Union Development and Structural Funds, Fondation FMC VIA-LR, RNSA

Conflict of interest:

J Bousquet reports personal fees and others from Chiesi, Cipla, Hikma, Menarini, Mundipharma, Mylan, Novartis, Sanofi-Aventis, Takeda, Teva, Uriach, others from Kyomed, outside the submitted work.

S Bosnic-Anticevich reports grants from TEVA, personal fees from TEVA, Boehringer Ingelheim, Sanofi, GSK, AstraZeneca, outside the submitted work.

E Correia-de-Sousa reports other from Boehringer Ingelheim, Novartis, grants from AstraZeneca, during the conduct of the study.

Dr. Cruz reports personal fees and non-financial support from Boehringer Ingelheim, AstraZeneca grants and personal fees from GSK, other from MSD, other from Sanofi, personal fees from Novartis, CHIESI, Eurofarma, Boston Scientific, outside the submitted work.

P Devillier reports personal fees from Astra Zeneca, GlaxoSmithKline, Meda Pharma, Sanofi, Novartis, Chiesi, outside the submitted work.

T Haahtela reports personal fees from Mundipharma, Novartis, and Orion Pharma, outside the submitted work.

JC Ivancevich reports personal fees from Faes Farma, EuroFarma Argentina, other from Sanofi Argentina, Lab. Casasco outside the submitted work.

J Just reports grants and personal fees from Novartis, ALK, Astra Zenaca, personal fees from Thermofischer, Zambon, outside the submitted work.

P Kuna reports personal fees from Adamed, Boehringer Ingelheim, AstraZeneca, personal fees from Chiesi, FAES, Berlin Chemie, Novartis, Polpharma, Allergopharma, outside the submitted work.

Dr. Kritikos reports personal fees from AstraZeneca, GlaxoSmithKline, Pfizer, outside the submitted work.

V Kvedariene has received payment for consultancy from GSK and for lectures from StallergensGreer, Berlin-Chemie outside the submitted work.

D Larenas Linnemann reports personal fees from GSK, Astrazeneca, MEDA, Boehringer Ingelheim, Novartis, Grunenthal, UCB, Amstrong, Siegfried, DBV Technologies, MSD, Pfizer. grants from Sanofi, Astrazeneca, Novartis, UCB, GSK, TEVA, Chiesi, Boehringer Ingelheim, outside the submitted work.

R Mösges reports personal fees from ALK, allergopharma, Allergy Therapeutics, Friulchem, Hexal, Servier, Klosterfrau, Bayer, FAES, GSK, MSD, Johnson&Johnson, Meda, Stada, UCB, Nuvo, grants from ASIT biotech, Leti, Optima, BitopAG, Hulka, Ursapharm, grants and personal fees from Bencard, Stallergenes, personal fees and non-financial support from Lofarma, Novartis, non-financial support from Atmos , Roxall, Bionorica, Otonomy, Ferrero, outside the submitted work.

Dr. Mullol reports personal fees from SANOFI-Genzyme-Regeneron, ALK-Abelló A/S, Menarini Group, MSD, GlaxoSmithKline, Novartis, UCB Pharma, GENENTECH – Roche, grants and personal fees from MYLAN-MEDA Pharma, URIACH Group, outside the submitted work.

R Naclerio : Advisory Boards: Sanofi, Novartis, ActoBio, Revance, Biomedical

Y Okamoto reports personal fees from Shionogi Co. Ltd., Torii Co. Ltd., GSK, MSD , Kyowa Co. Ltd., from Eizai Co. Ltd., grants and personal fees from Kyorin Co. Ltd., Tiho Co. Ltd., grants from Yakuruto Co. Ltd., Yamada Bee Farm, outside the submitted work.

N Papadopoulos reports personal fees from Novartis, Faes Farma, BIOMAY, HAL, Nutricia Research, Menarini, Novartis, MEDA, Abbvie, Novartis, MSD, Omega Pharma Danone, grants from Menarini outside the submitted work.

O Pfaar reports grants and personal fees from ALK-Abelló, Allergopharma, Stallergenes Greer, HAL Allergy Holding B.V./HAL Allergie GmbH, Bencard Allergie GmbH/Allergy Therapeutics, Lofarma, Biotech Tools S.A, LETI/LETI Pharma, Anergis S.A. grants from Biomay, Nuvo, Circassia, Glaxo Smith Kline, personal fees from Novartis Pharma, MEDA Pharma, Mobile Chamber Experts (a GA²LEN Partner), Pohl-Boskamp, Indoor Biotechnologies, grants from, outside the submitted work.

D Plavec reports grants and personal fees from GlaxoSmithKline, personal fees from Menarini, Pliva, AbbVie, Novartis, MSD, Chiesi, Revenio personal fees and non-financial support from Boehringer Ingelheim, non-financial support from Philips, outside the submitted work.

This article is protected by copyright. All rights reserved.

A Todo-Bom reports grants and personal fees from Novartis, Boehringer Ingelheim, Mundipharma, GSK (GlaxoSmithKline), Teva Pharma, AstraZeneca, grants from Leti, outside the submitted work.

M Wagenmann reports personal fees from AstraZeneca, HAL Allergy, MEDA Pharma, Stallergenes, ALK-Abelló, Teva, grants and personal fees from Allergopharma, Sanofi-Aventis, , grants from Allakos, grants from F. Hoffmann-La Roche, GlaxoSmithKline, Otonomy, Strekin, outside the submitted work.

S Wasserman reports personal fees from Merck, GSK, Novartis, Behring, Shire, Sanofi, Barid Aralez, Mylan Meda, Pediapharm outside the submitted work.

T Zuberbier reports and Organizational affiliations: Committee member: WHO-Initiative "Allergic Rhinitis and Its Impact on Asthma" (ARIA), Member of the Board: German Society for Allergy and Clinical Immunology (DGAKI)

Head: European Centre for Allergy Research Foundation (ECARF) , Secretary General: Global Allergy and Asthma European Network (GA²LEN) ,Member: Committee on Allergy Diagnosis and Molecular Allergology, World Allergy Organization (WAO).

Statement of Contribution :

S Bosnic-Anticevich (1) and J Bousquet planned the project and wrote the first draft of the paper.

E Costa (2), E Menditto (3), O Lourenço (4), E Novellino (5), S Bialek (6), V Briedis (7), R Buonaiuto (8), H Chrystyn (9), B Cvetkosvki (1), S Di Capua (10), V Kritikos (1), A Mair (11), V Orlando (3), E Paulino (12), J Salimäki (13), R Söderlund (14), R Tan (1), DM Williams (15), P Wroczynski (6) were pharmacists who reviewed the paper and made some additive concepts.

I Agache (16), IJ Ansotegui (17), JM Anto (18-21), A Bedbrook (22), C Bachert (23), M Bewick (24), C Bindeslev-Jensen (25), J Brozek (26), GW Canonica (27), V Cardona (28), W Carr (29), T Casale (30), NH Chavannes (31), J Correia de Sousa (32), AA Cruz (33), W Czarlewski (34), G De Carlo (35), P Demoly (36, 37), P Devillier (38), MS Dykewicz (39), M Gaga (40), Y El-Gamal (41), J Fonseca (42), WJ Fokkens (43), MA Guzmán (44), T Haahtela (45), PW Hellings (46), M Illario (47), JC Ivancevich (48), J Just (49), I Kaidashev (50), M Khaitov (51), N Khaltayev (52), T Keil (53), L Klimek (54), ML Kowalski (55), P Kuna (56), V Kvedariene (57), D Larenas-Linnemann (58), D Laune (59), LTT Le (60), KC Lodrup Carlsen (61), B Mahboub (62), D Maier (63), J Malva (64), P Manning (65), M Morais-Almeida (66) , R Mösges (67), J Mullol (68) , L Münter (69), R Murray (70), R Naclerio (71), L Namazova-Baranova (72), K Nekam (73), TD Nyembue (74), K Okubo (75), RE O'Hehir (76), K Ohta (77), Y Okamoto (78), GL Onorato (22), S Palkonen (35), P Panzner (79), NG Papadopoulos (80), HS Park (81), R Pawankar

This article is protected by copyright. All rights reserved.

(82), O Pfaar (83), J Phillips (84), D Plavec (85), TA Popov (86), P Potter (87), EP Prokopakis (88), RE Roller-Wirnsberger (89), M Rottem (90), D Ryan (91), B Samolinski (92), M Sanchez-Borges (93), HJ Schunemann (26), A Sheikh (94), JC Sisul (95), D Somekh (96), C Stellato (97), T To (98), A Todo-Bom (99), PV Tomazic (100), S Toppila-Salmi (45), A Valero (101), A Valiulis (102), E Valovirta (103), MT Ventura (104), M Wagenmann (105), D Wallace (106), S Wasserman (107), M Wickman (108), PK Yiallourous (109), A Yorgancioglu (110), OM Yusuf (111), HJ Zar (112), ME Zernotti (113), L Zhang (114), M Zidarn (115), T Zuberbier (116), are members of the ARIA-MASK group. They all participated (i) to the concept of the paper, (ii) the revision of the draft and (iii) will deploy of ARIA in the pharmacy in their own country. ARIA in the pharmacist will be used as a multistakeholder care pathway and a multidisciplinary group is needed.

All authors approved the paper

Abstract

Pharmacists are trusted health professionals. Many patients use over-the-counter (OTC) medications and are seen by pharmacists who are the initial point of contact of allergic rhinitis management in most countries. The role of pharmacists in integrated care pathways (ICPs) for allergic diseases is important. This paper builds on existing studies and provides tools intended to help pharmacists provide optimal advice/interventions/strategies to patients with rhinitis. The ARIA-pharmacy ICP includes a diagnostic questionnaire specifically focusing attention on key symptoms and markers of the disease, a systematic Diagnosis Guide (including differential diagnoses) and a simple flowchart with proposed treatment for rhinitis and asthma multimorbidity. Key prompts for referral within the ICP are included. The use of technology is critical to enhance the management of AR. However, the ARIA-pharmacy ICP should be adapted to local health care environments/situations as regional (national) differences exist in pharmacy care.

Abbreviations

AHA: Active and healthy ageing

AIRWAYS ICPs: Integrated care pathways for airway diseases

AR: Allergic rhinitis

ARIA: Allergic Rhinitis and Its Impact on Asthma

This article is protected by copyright. All rights reserved.

CARAT: Control of Allergic Rhinitis and Asthma Test

CDSS: Clinical decision support system

CRD: Chronic Respiratory Diseases

DG CONNECT: Directorate General for Communications Networks, Content and Technology

DG Santé: Directorate-General for Health and Food Safety

DG: Directorate General

EFA: European Federation of Allergy and Airways Diseases Patients' Associations

EIP on AHA: European Innovation Partnership on AHA

EIP: European Innovation Partnership

GARD: WHO Global Alliance against Chronic Respiratory Diseases

HCP: Health care professional

ICP: Integrated care pathway

JA-CHRODIS: Joint Action on Chronic Diseases and Promoting Healthy Ageing across the Life Cycle

MACVIA: contre les MALadies Chroniques pour un Vieillissement Actif (Fighting chronic diseases for AHA)

MASK: Mobile Airways Sentinel Network

MedALL: Mechanisms of the Development of ALLergy (FP7)

mHealth: Mobile health

NCD: Non-communicable disease

OTC: Over-the-counter

POLLAR: Impact of air POLLution on Asthma and Rhinitis

QOL: Quality of life

SCUAD: Severe chronic upper airway disease

VAS: Visual analogue scale

WHO: World Health Organization

WPAI-AS: Work Productivity and Activity questionnaire

This article is protected by copyright. All rights reserved.

Key words: ARIA, asthma, care pathways, pharmacist, rhinitis

Introduction

Allergic diseases such as rhinitis and asthma are common, complex conditions, associated with allergen-specific IgE and non-allergic mechanisms (1, 2). These diseases represent an enormous burden associated to personal, medical, and social costs as well as impairment in work productivity (3-6), impacting health and social inequalities in all age groups (7).

Allergic rhinitis (AR) is a highly diverse chronic disease spanning from mild intermittent rhinitis to Severe Chronic Upper Airway Disease (SCUAD) (8). There is evidence that the condition is sub-optimally managed with several unmet needs which include:

- Understanding the different endotypes (rhinitis and asthma) (9) and the presence of different phenotypes of AR and multi-morbidities which can impact AR and asthma control,
- Improving AR diagnosis using modern technology (10),
- Delivering management strategies and interventions which address suboptimal rhinitis and asthma control arising not only from suboptimal medical management/treatment, but as a result of a range of cultural and/or social barriers (8, 11),
- Assessing risk factors such as allergens and pollutants to incorporate them into management strategies, using technologies that promote sentinel networks within multidisciplinary care pathways (i.e. integrated care pathways (ICPs)) (12),
- Stratifying patients, based on their needs, in order to optimize the use and effectiveness of ICPs (13), and
- Promoting multidisciplinary teams within integrated ICPs, endorsing innovation in clinical trials and encouraging patient empowerment.

ICPs are structured multidisciplinary care plans which detail essential steps in the care of patients. They promote the translation of guidelines into local protocols and their subsequent application to clinical practice. They empower patients and their carers (health and social). ICPs differ from practice guidelines as they are utilized by a multidisciplinary team and have a focus on the quality and

coordination of care. Pharmacists are at the forefront of ICPs for AR. An ICP is intended to act as a guide to treatment.

This paper builds on existing studies and provides tools intended to help pharmacists provide optimal advice/interventions/strategies to patients with AR. The ARIA-Pharmacy ICP includes a diagnostic questionnaire specifically focusing attention on key symptoms and markers of the disease, a systematic Diagnosis Guide (including a differential diagnosis) and a simple flowchart with proposed treatment for rhinitis and asthma multimorbidity. Key prompts for referral within the ICP are included. The use of technology is critical to enhance the management of AR. However, the ARIA-pharmacy ICP should be adapted to local health care environments/situations.

1- Pharmacist challenge in allergic rhinitis

Considering the challenges associated with AR and the identified needs, it is clear that the pharmacist has a role for AR management in practice, through a guided change management process. This is both within the scope of pharmacy practice and in line with future models of integrated care. Pharmacists are ideally placed to manage this extremely important link in care pathways.

Worldwide, pharmacists receive advanced training in basic and clinical sciences. Given the importance of self-medication in many allergic diseases and in iatrogenic disease, pharmacist interventions are well placed to maximize the benefits and minimize the adverse events associated with pharmacotherapy. This is particularly important in AR and asthma multi-morbidity (14) as well as in the elderly patients, a large number of whom present with allergic diseases in combination with other chronic diseases (15). Moreover, most AR medications are available over the counter (OTC) (16-19). The impact of the switch from prescription to OTC medications has been profound in AR (20) with a significant impact on cost and health utilization reduction (21). Therefore, as trusted healthcare professionals in the community, pharmacists are well placed to play a critical role identifying the symptoms of AR, recommending appropriate OTC treatment (22-24) and integrating ICPs into health care teams.

Pharmacists (along with patients, clinicians and other healthcare professionals (HCPs)) are faced with the relative merits and downsides of the various treatment options. Clinical practice guidelines for AR management developed over the past 20 years have improved the care of AR patients (25) and provide a critical framework for AR management (25-27). These guidelines are becoming particularly

important for HCPs in primary care since, in most countries, few AR patients consult a specialist physician. In fact, many patients with AR under-recognize their condition (4) and, as such, do not even consult a physician (28). A large proportion of AR patients actually self-manage their condition (5, 6) and the pharmacist is often the first HCP to whom a person with nasal symptoms presents (29, 30). This further strengthens the important role of pharmacists within multidisciplinary health care teams, acting at different steps of the ARIA-Pharmacy ICPs (23, 31-33).

The specific role of pharmacists within ICPs can be evidenced through several strategies that have been initiated (34) or completed in the AR management in the pharmacy, and in studies confirming the important impact of pharmacist interventions on AR outcomes (18, 35-43). The specific areas of pharmacist impact in AR are summarized in Table 1.

2- ARIA

Allergic Rhinitis and its Impact on Asthma (ARIA) commenced during a World Health Organization (WHO) workshop in 1999 (44), and was further developed by the WHO Collaborative Center for Rhinitis and Asthma (2002-2013). This work has been undertaken in four phases, which are briefly summarized below:

- Phase 1: The initial goals were (i) to propose a new AR classification, (ii) to promote the concept of multi-morbidity in asthma and rhinitis and (iii) to develop guidelines with all stakeholders that could be used globally for all countries and populations. ARIA has been disseminated and implemented in over 70 countries globally (7, 45-54), and was revised and updated in 2008 (7).
- Phase 2: Focused on transparent reporting of guidelines to facilitate understanding and acceptance using the GRADE approach (25, 26).
- Phase 3: MASK (Mobile Airways Sentinel Network), an ARIA initiative, is focusing on (i) the implementation of multi-sectoral ICPs (ii) using emerging technologies (iii) with real-world data (iv) for individualized and predictive medicine (v) in rhinitis and asthma multi-morbidity, (vi) by a multi-disciplinary group or by patients themselves (self-care) using the AIRWAYS ICPs algorithm (Figure 2) (vii) across the life cycle (12, 55).
- Phase 4 concerns change management strategies (56). ARIA-pharmacy ICPs represent one of the change management steps.

ARIA in the pharmacy (23) and its pocket guide were published in 2004 to help pharmacists with the management of AR symptoms and their impact on asthma. Most recommendations proposed in

2004 are still valid but should be updated due to the larger number of medications now available OTC. In the initial approach, ICPs were not considered and the new information and communication technology (ICT) was not available.

3- Integrated care pathways for allergic rhinitis

a. AIRWAYS ICPs

The B3 Action Plan of the European Innovation Partnership on Active and Healthy Ageing (EIP on AHA, DG Santé and DG CONNECT) (57) promotes integrated care models for chronic diseases, including the use of remote monitoring. Based on this initiative and its aims to produce evidence-based translational outcomes, an ICP for airways disease (AIRWAYS ICPs) was developed. Its aim was to launch a collaboration to develop practical multi-sectoral ICPs (i) to reduce chronic respiratory disease (CRD) burden, mortality and multi-morbidity, (ii) to improve education of all stakeholders, (iii) to improve work productivity, (iv) to promote AHA and (v) to reduce inequities in all populations globally (58).

AIRWAYS ICPs considers a multi-disciplinary approach to AR and asthma multi-morbidity management with the pharmacist at the forefront of the algorithm (Figure 1). A very large number of AR patients use OTC drugs (23) and are treated in community pharmacies while the vast majority of patients who visit primary care physicians or specialists have moderate/severe rhinitis (59-63).

However, pharmacy practice varies widely across countries and ICPs should be tailored to the local needs taking into account cultural barriers, socio-economic considerations, health care practices and available OTC medications.

b. mHealth in the management of AR incorporating technology for better AR management

mHealth including apps running on consumer smart devices (i.e. smartphones and tablets) has the potential to profoundly impact healthcare (64). An evidence-based app for use in AR management is available. MASK (Mobile Airways Sentinel network) (ARIA Phase 3) is an app, i.e. an information and communications technology (ICT) system, centred around the patient (12, 55, 65). It is an implementation tool of the European Innovation Partnership on Active and Healthy Ageing (EIP on AHA (57, 66)). A mobile phone App (available for Android and iOS), the *Allergy Diary*, launched in 23

This article is protected by copyright. All rights reserved.

countries and 17 languages (55, 67), is associated with an inter-operable tablet for physicians and other HCPs (65), using visual analogue scale (VAS), the same simple common language to manage AR ((68, 69).

The *Allergy Diary* also encompasses questions relating to the patient's experience of AR symptoms, an additional quality of life questionnaire (EQ-5D (EuroQuol)) (70-72), and a validated tool for simultaneously assessing AR and asthma control (Control of Allergic Rhinitis and Asthma Test (CARAT)) (73-76). These can be used for self-monitoring or can be used for inclusion into clinical trials. The *Allergy Diary* is an integrated approach and can be used by patients in the management of their AR, by HCPs in assisting and guiding management and by researchers to collect large-scale population data on AR status and management globally.

c. Clinical decision support system: additional resource for pharmacists

Clinical decision support systems (CDSS) are software algorithms that advise HCPs on diagnosis and management of patients based on the interaction of patient data and medical information, such as prescribed drugs. CDSS should be based on the best evidence to aid patients and HCPs in shared decision making. In a prospective intervention study, a computerized pharmacy CDSS for the counselling of patients with AR was tested. The results showed that pharmacists omitted many questions mandatory to assessing whether self-medication is appropriate, showing the importance of the CDSS (77).

An AR CDSS (MASK CDSS) is available for pharmacists as a companion to the Allergy Diary (*ARIA Allergy Diary Companion*) (12, 55). The *ARIA Allergy Diary Companion* is based on an algorithm to aid in the selection of pharmacotherapy for patients with AR and to stratify their disease severity (55). It uses a simple step-up/step-down individualized approach to AR pharmacotherapy and may hold the potential for optimal control of symptoms, while minimizing side-effects and costs. Its use is encouraged for pharmacists, however its application may vary depending on the availability of medications in the different countries and on resources.

4- Management of allergic rhinitis in the pharmacy: ARIA-pharmacy ICP

Based on the ICP concept, mHealth and CDSS, ARIA Phase 4 has developed an ICP for Community Pharmacy, to assist pharmacists in the management of AR within the AIRWAY ICPs framework (Figure 1). In summary, this frames the role of the pharmacist in AR across 4 domains:

This article is protected by copyright. All rights reserved.

- Recognition and classification of allergic rhinitis symptoms. As pharmacists are often the health care location at which people with AR either initially present or most frequently present, recognizing and classifying AR becomes critical.
- Identification of AR-related ocular symptoms and asthma: Pharmacist's role includes not only assisting in confirming that AR is present but will allow for the identification of possible other conditions and patients at risk, requiring immediate referral to a general practitioner.
- AR treatment: Initiating/recommending optimal treatment, as medication management has been proven to be the most effective way of managing AR (25, 26).
- Patient support and AR monitoring over time: Supporting the patient with appropriate evidence-based education, self-management support and long term monitoring over time.

These 4 domains are discussed below:

a. Recognition and classification of allergic rhinitis symptoms

Although AR diagnosis requires tests to confirm the allergic sensitization, most patients self-diagnose their AR, consulting pharmacists without a doctor diagnosis of AR. Several questionnaires are available for the screening of allergic diseases (78, 79). Most AR patients have multiple nasal symptoms (rhinorrhoea, sneezing, nasal pruritus and obstruction), a large percentage have ocular symptoms (ocular tearing, redness and pruritus) and many have asthma. In order for pharmacists to evaluate the possibility of screening for AR, a series of questions should be asked to patients with nasal symptoms. The questionnaire in Table 2, while it will not be able to lead to a definitive diagnosis of AR, may enable pharmacists to determine whether a diagnosis of AR should be further investigated and can assist in identifying warning symptoms that need further medical investigation.

Figure 2 summarises the list of symptoms, which, if they occur in isolation, may be suggestive of a condition other than AR. The key feature of AR to remember is that AR symptoms are never unilateral and minimal bleeding may occur during long-term intranasal therapy. Furthermore, although nasal obstruction, loss of smell (80), facial pain or post nasal drip may be AR symptoms, when they occur as single symptoms, they are unlikely to be from allergic origin. Purulent discharge, especially if accompanied with fever, is suggestive of an infection (Figure 2). AR may present with symptoms similar to those of a number of other conditions induced by non-allergic triggers including viral infections such as the common cold. Table 2 may help the pharmacist to differentiate allergy from other causes including infection.

The ARIA guideline proposes a classification of AR based on symptom control, QOL and daily impact as well as duration (7, 44). Disease control associated with several health outcomes, including QOL (59-63) and some mHealth endpoints (72, 81-84), should be considered in ICPs. Although the duration of rhinitis is an important indicator of asthma multimorbidity (85), and duration and efficacy of treatment in AR (61), AR control should be considered as the most important end point for the pharmacist.

AR 'control' in AR patients is the main goal of treatment (86). Control is the degree to which therapy goals are currently met such as glycaemic control in diabetes. However, measures of AR control are somewhat different to many other chronic diseases as while they include symptom scores, alternative scales/scores, which are reflective of the impact of AR on day-to-day living, have been shown to be validated as measure of AR control. For example visual analogue scales (VAS) assessing impact of AR (69, 87), quality of life (QOL) measures and scores with several items which cover symptoms and impact on daily living (88, 89) are commonly used. In deciding which measure of AR control is most appropriate, it is important for a score to be simple and responsive to change over time, as AR control changes, hence VAS scores, which can be used in all age groups, (90, 91) are available in a wide variety of languages (92, 93) and have been shown to be valid for assessing AR control, are useful instruments for assessing AR control (Figure 3).

A VAS (range: 0-20) can be used to assess control before and some days after treatment (Figure 6). If the scale is <20/100, the patient has controlled AR, if the scale is from 20 to 50/100, the patient has partly controlled disease and for $\geq 50/100$, the patient has uncontrolled disease (Figure 6). Using VAS can help the pharmacist to assess response to OTC treatment. VAS can be used to assess overall AR control (VAS global measure) as it relates to nasal symptoms and eye symptoms (94) (Figure 3 and Figure 6). There is a high degree of correlation between VAS global measured ("Overall how much are your allergic symptoms bothering you today?") and VAS for nose symptoms (81, 84). An electronic form of the VAS exists in the *Allergy Diary* (81, 84) and has been validated against several end points.

b. Identification of AR-related ocular symptoms and asthma

Ocular symptoms are commonly associated with nasal symptoms in AR, and they can be diagnosed using simple questions. However, some forms of conjunctivitis require referral to a physician (Figure 4). A VAS for ocular symptoms may also be used and is included in the *Allergy Diary* (Figure 3).

Asthma is a common multi-morbidity of AR and should be checked. A proposal has been made in ARIA in the pharmacy (23) (Figure 5). All patients with rhinitis should be evaluated for asthma, particularly if they have persistent and/or moderate-severe rhinitis.

c. AR Treatment

Goals for the treatment of rhinitis should be determined after an accurate diagnosis of AR and a validated assessment of control, including evaluation of coexisting asthma. AR treatment goals include:

- Normal sleep.
- Ability to undertake normal daily activities, including work and school attendance, without limitation or impairment, and the ability to participate fully in sport and leisure activities.
- No troublesome symptoms.
- No or minimal side-effects of rhinitis treatment. It should be recognized that many OTC drugs for the treatment of AR can induce sedation and should be avoided.

In all guidelines, it has been considered that many medications may be used at several steps of severity and duration (44); including antihistamines (oral and intranasal), intranasal glucocorticosteroids (INCS) leukotriene receptor antagonists and chromones (intranasal and eye drops). Anticholinergics and decongestants (intranasal and oral) are sometimes noted in the treatment of AR on a short-term basis however, there is little good quality evidence for their effectiveness (45) and prolonged use of decongestants (>10 days) may lead to rebound swelling of the nasal mucosa, drug-induced rhinitis (also known as rhinitis medicamentosa) and tachyphylaxis (7). Table 3 summarises the specific pharmacological effects of these different agents on specific AR symptoms (95).

In consideration of the specific role of the pharmacist, however, it is important to focus on pharmacotherapy that is available OTC in pharmacies i.e. antihistamines and in some countries, intranasal corticosteroids. Considering OTC medications, intra-nasal corticosteroids (INCS) are more effective than oral or intranasal H1-antihistamines (27, 44) but many patients prefer oral drugs. Other medications include leukotriene antagonists, nasal washing, vasoconstrictors and cromoglycate. In some countries (currently in New Zealand), the combination of azelastine and

fluticasone propionate is OTC. This medication may be preferred if the patient wants a rapid onset of action of the treatment (26).

Figure 6 provides a clinical treatment pathway to help pharmacists choose the appropriate OTC medications, based on assessment of AR control using VAS. The patient's preference should always be considered. OTC medication availability differs between countries.

Immunotherapy and AR treatment: The application of both subcutaneous immunotherapy and sublingual immunotherapy (for treatment of allergies to pollens, ragweed and house dust mite) are reserved for patients with severe AR. They have been shown to improve the quality of life of people with AR and reduce the need for other pharmacological therapy (96-98). Currently in most countries, initiation of immunotherapy is both administered and monitored by specialist physicians. The role of pharmacy within the context of this treatment has not as yet been defined.

Non pharmacological AR treatment: In addition to pharmacological treatment, historically patients with AR have been recommended non-pharmacological strategies such as allergen avoidance or minimisation and nasal washing. When it comes to seasonal AR, it is evident that avoidance of seasonal allergens is effective. i.e. patients with seasonal AR do not experience symptoms outside of season. However, while the level of evidence is low, during pollen season, patients can be advised to close windows at night, drive with closed windows and wear wrap-around glasses when outdoors, to prevent exacerbation of symptoms; and wear sunglasses (99), nasal filters (100) and apply balms and ointments to the nose (95) during pollen season to reduce symptoms. For patients with occupational AR, avoiding exposure to the occupational agent trigger AR is recommended (101).

There is limited evidence that saline washings, irrigation or sprays are effective in reducing AR symptoms and potentially reducing the amount of pharmacotherapy needed (7, 102) however, they are well tolerated, safe and inexpensive(103), hence for patients who would like to try them, they can be safely recommended.

d. Patient Support and AR monitoring over time

In considering the way in which pharmacists can support the patient with AR, it is important to build on the process of recognition, classification and treatment recommendations with appropriate education, self-management support and both short and long term monitoring.

Education around the condition of AR is fundamental, however, education about the way in which treatment works, the need to be both adherent and use intranasal devices correctly, is also critical. In particular, it is well recognized that non-adherence to AR treatment adds to the burden of disease. It is difficult to accurately determine non-adherence to AR treatment, as some medications are taken seasonally (as needed) and many are available over the counter in the pharmacy. It is, however estimated that for people using intranasal corticosteroids to treat their AR, adherence is approximately 35% (104). Supporting the patient to better understand the way in which medication works and the need to take it regularly is critical. It has been shown that when pharmacists work with patients to assist them in setting their goals for AR management, better long term AR outcomes are achieved (36, 37)

Further to this, patients must be shown how to use their intranasal sprays correctly. Little is known about the proportion of patients who use their intranasal medication correctly however, the way in which the patient uses their intranasal spray does impact on way in which the spray distributes in the nasal cavity (105) and the occurrence of local side effects (106). Therefore, as with asthma devices, it is recommended that pharmacists train the patient how to use their intranasal device with a placebo device.

Monitoring adherence with AR treatment is a challenge. This is where the use of technology can assist. There are several mHealth tools for AR follow up but only the *Allergy Diary* has been validated. The Allergy Diary (MASK-rhinitis) has been implemented in 23 countries and 16 languages. Over 23,000 users have been recorded (55, 81, 82, 107-111). This tool appears to be appropriate for pharmacists, in particular since it is extremely simple, does not require the pharmacist to undertake any particular education on its use, is user friendly for the patient and can be used for follow-up over time.

For confidentiality reasons, patients cannot give access of the electronic data that they record in the *Allergy Diary* (both symptom VAS and medication use) to HCPs. However, they can print their daily AR control VAS responses and medication use, as summarized in Figure 7. Comparison of control at first dispensing of an OTC medication with the evolution of control during treatment will guide pharmacists to stop or increase the treatment with OTC medications or suggest referral to a physician. Moreover, this control chart will help the physician to optimize the treatment.

Conclusion

The present paper has summarized the need for community pharmacists to play an important role in ICPs for AR. Although each recommendation varies between countries due to the available OTC medications, legislation and cultural differences, ARIA-Pharmacy may be used as a model for the implementation of ICPs in different countries.

Table 1: Importance of pharmacists in the management of allergic rhinitis

- | |
|---|
| <ul style="list-style-type: none">• Recognizing (identification of) AR• Risk assessment/stratification• Over-the-counter treatment• Patient education• Referral to a physician• Administration of topical treatment technique, including teaching and re-evaluation of treatment technique• Adherence to treatment• Research on database |
|---|


Table 2: Is it a cold or allergic rhinitis?

| Symptoms | Cold | Allergic rhinitis |
|---------------------------|---|---|
| Duration | 5-10 days | Variable but can last months or present recurrent episodes |
| Season | Most often in winter but may be possible at any time | Any time of the year, more common during pollen seasons which can occur in winter |
| Course of the disease | Symptoms usually take a few hours or days to be severe | Symptoms can be severe within minutes |
| Nasal and ocular symptoms | <ul style="list-style-type: none"> • Severe nasal obstruction • Rhinorrhoea • Sneezing (rarely in bouts) • NO nasal or ocular pruritus • Rare ocular symptoms • Frequent loss of smell | <ul style="list-style-type: none"> • Profuse watery rhinorrhoea • Several episodes of sneezing in succession • Nasal pruritus • Variable nasal congestion • Often associated with ocular symptoms (tearing, redness, pruritus) • Partial loss of smell in the most severe patients |
| Sore Throat | Common | Sometimes |
| Cough | Common | Sometimes |
| Chest Discomfort | <ul style="list-style-type: none"> • Mild to moderate • Cold can lead to severe asthma exacerbation | Rare, except for those with allergic asthma |

Table 3: AR symptoms and relative effectiveness of different pharmacotherapeutic agents (from (95, 112))

| | Sneezing | Rhinorrhea | Nasal obstruction | Nasal itch | Eye symptoms |
|---|----------|------------|-------------------|------------|--------------|
| Hi-antihistamines | | | | | |
| Oral | ++ | ++ | + | +++ | ++ |
| Intranasal | ++ | ++ | + | ++ | 0 |
| Eye drops | 0 | 0 | 0 | 0 | +++ |
| Corticosteroids | | | | | |
| Intranasal | +++ | +++ | ++ | ++ | ++ |
| Chromones | | | | | |
| Intranasal | + | + | + | + | 0 |
| Eye drops | 0 | 0 | 0 | 0 | + |
| Decongestants | | | | | |
| Intranasal | 0 | 0 | ++++ | 0 | 0 |
| Oral | 0 | 0 | + | 0 | 0 |
| Anti-cholinergics | 0 | | | 0 | 0 |
| Anti-leukotrienes | 0 | | | 0 | ++ |
| Intranasal steroids and intranasal antihistamine 1 | +++ | +++ | +++ | +++ | +++ |

Figure 1: ARIA in the pharmacy initiative for AIRWAYS ICPs in allergic rhinitis (modified from (66))



This is a generic algorithm which varies between countries depending on regulations.

The health care professional should ascertain that the treatment taken by the patient accords to best practices. In particular, regular use of short-acting β_2 agonist as a single treatment should be avoided. Similarly, prolonged use of intra-nasal vasoconstrictors in rhinitis should be avoided.

Figure 2: Diagnosis of allergic rhinitis in the pharmacy (Adapted from (23))




Figure 3: The MASK Allergy Diary




Figure 4: Diagnosis of allergic conjunctivitis at the pharmacy (adapted from (23))




Figure 5: Screening of asthma in rhinitis patients in the pharmacy (from (23))





Figure 6: Treatment of allergic rhinitis in the pharmacy (adapted from (23))



VAS nose: "How are your nose symptoms bothering you today?"

*INCS if coexisting asthma.

AH antihistamine; INCS intranasal corticosteroid; INAH intranasal antihistamine.

VAS eyes: "How are your eye symptoms bothering you today?"


*INCS if coexisting asthma.

IOAH intraocular antihistamine; INCS intranasal corticosteroid.

This algorithm should be adapted to the regulations, needs, price of medications and cultural barriers of each country (or region)

Figure 7: Transfer of patient information on a computer and printed information (From (109))

- 1- Open the Allergy Diary app and choose "Show Data on Computer" in the main menu
- 2- Go to www.macvia-aria-allergy-diary.com/data on your PC/Laptop (enter this URL in the address bar of the browser from your PC/Laptop)
- 3- Scan the QR code with the Allergy Diary app
- 4- The screen with your personal data can be seen
- 5- And you can also print these data (see figure below)



References

1. Bousquet J, Anto J, Auffray C, Akdis M, Cambon-Thomsen A, Keil T, et al. MeDALL (Mechanisms of the Development of ALLergy): an integrated approach from phenotypes to systems medicine. *Allergy*. 2011;66(5):596-604.
2. Cingi C, Gevaert P, Mosges R, Rondon C, Hox V, Rudenko M, et al. Multi-morbidities of allergic rhinitis in adults: European Academy of Allergy and Clinical Immunology Task Force Report. *Clin Transl Allergy*. 2017;7:17.
3. Vandenplas O, Vinnikov D, Blanc PD, Agache I, Bachert C, Bewick M, et al. Impact of Rhinitis on Work Productivity: A Systematic Review. *J Allergy Clin Immunol Pract*. 2018;6(4):1274-1286..
4. Cvetkovski B, Kritikos V, Yan K, Bosnic-Anticevich S. Tell me about your hay fever: a qualitative investigation of allergic rhinitis management from the perspective of the patient. *NPJ Prim Care Respir Med*. 2018;28(1):3.
5. Tan R, Cvetkovski B, Kritikos V, Price D, Yan K, Smith P, et al. Identifying the hidden burden of allergic rhinitis (AR) in community pharmacy: a global phenomenon. *Asthma Res Pract*. 2017;3:8.
6. Tan R, Cvetkovski B, Kritikos V, Price D, Yan K, Smith P, et al. The Burden of Rhinitis and the Impact of Medication Management within the Community Pharmacy Setting. *J Allergy Clin Immunol Pract*. 2018;6(5):1717-1725
7. Bousquet J, Khaltaev N, Cruz AA, Denburg J, Fokkens WJ, Togias A, et al. Allergic Rhinitis and its Impact on Asthma (ARIA) 2008 update (in collaboration with the World Health Organization, GA(2)LEN and AllerGen). *Allergy*. 2008;63 Suppl 86:8-160.
8. Bousquet J, Bachert C, Canonica GW, Casale TB, Cruz AA, Lockey RJ, et al. Unmet needs in severe chronic upper airway disease (SCUAD). *J Allergy Clin Immunol*. 2009;124(3):428-33.
9. De Greve G, Hellings PW, Fokkens WJ, Pugin B, Steelant B, Seys SF. Endotype-driven treatment in chronic upper airway diseases. *Clin Transl Allergy*. 2017;7:22.
10. Alessandri C, Ferrara R, Bernardi ML, Zennaro D, Tuppo L, Giangrieco I, et al. Diagnosing allergic sensitizations in the third millennium: why clinicians should know allergen molecule structures. *Clin Transl Allergy*. 2017;7:21.
11. Bousquet J, Mantzouranis E, Cruz AA, Ait-Khaled N, Baena-Cagnani CE, Bleecker ER, et al. Uniform definition of asthma severity, control, and exacerbations: document presented for the World Health Organization Consultation on Severe Asthma. *J Allergy Clin Immunol*. 2010;126(5):926-38.
12. Bousquet J, Schunemann HJ, Fonseca J, Samolinski B, Bachert C, Canonica GW, et al. MACVIA-ARIA Sentinel Network for allergic rhinitis (MASK-rhinitis): the new generation guideline implementation. *Allergy*. 2015;70(11):1372-92.
13. Hellings PW, Fokkens WJ, Bachert C, Akdis CA, Bieber T, Agache I, et al. Positioning the Principles of Precision Medicine in Care Pathways for Allergic Rhinitis and Chronic Rhinosinusitis - an EUFOREA-ARIA-EPOS-AIRWAYS ICP statement. *Allergy*. 2017;72(9):1297-305.

- Accepted Article
14. Emmerton LM, Smith L, LeMay KS, Krass I, Saini B, Bosnic-Anticevich SZ, et al. Experiences of community pharmacists involved in the delivery of a specialist asthma service in Australia. *BMC Health Serv Res.* 2012;12:164.
 15. Kouladjian L, Chen TF, Gnjdjic D, Hilmer SN. Education and Assessment of Pharmacists on the Use of the Drug Burden Index in Older Adults Using a Continuing Professional Development Education Method. *Am J Pharm Educ.* 2016;80(4):63.
 16. Sullivan PW, Nair KV, Patel BV. The effect of the Rx-to-OTC switch of loratadine and changes in prescription drug benefits on utilization and cost of therapy. *Am J Manag Care.* 2005;11(6):374-82.
 17. Rachelefsky G, Farrar JR. Are you comfortable with over-the-counter intranasal steroids for children? A call to action. *J Allergy Clin Immunol Pract.* 2014;2(3):271-4.
 18. Carney AS, Price DB, Smith PK, Harvey R, Kritikos V, Bosnic-Anticevich SZ, et al. Seasonal patterns of oral antihistamine and intranasal corticosteroid purchases from Australian community pharmacies: a retrospective observational study. *Pragmat Obs Res.* 2017;8:157-65.
 19. OTC fluticasone furoate nasal spray (Flonase Sensimist) for allergic rhinitis. *Med Lett Drugs Ther.* 2017;59(1519):e70-e1.
 20. Trygstad TK, Hansen RA, Wegner SE. Evaluation of product switching after a state Medicaid program began covering loratadine OTC 1 year after market availability. *J Manag Care Pharm.* 2006;12(2):108-20.
 21. Hay JW, Leahy M. Cost and utilization impacts of oral antihistamines in the California Medi-Cal program. *Value Health.* 2005;8(4):506-16.
 22. Fromer LM, Blaiss MS, Jacob-Nara JA, Long RM, Mannion KM, Lauersen LA. Current Allergic Rhinitis Experiences Survey (CARES): Consumers' awareness, attitudes and practices. *Allergy Asthma Proc.* 2014;35(4):307-15.
 23. ARIA in the pharmacy: management of allergic rhinitis symptoms in the pharmacy. *Allergic rhinitis and its impact on asthma.* 2004;59(4):373-87.
 24. Carr WW, Yawn BP. Management of allergic rhinitis in the era of effective over-the-counter treatments. *Postgrad Med.* 2017;129(6):572-80.
 25. Brozek JL, Bousquet J, Baena-Cagnani CE, Bonini S, Canonica GW, Casale TB, et al. Allergic Rhinitis and its Impact on Asthma (ARIA) guidelines: 2010 revision. *J Allergy Clin Immunol.* 2010;126(3):466-76.
 26. Brozek JL, Bousquet J, Agache I, Agarwal A, Bachert C, Bosnic-Anticevich S, et al. Allergic Rhinitis and its Impact on Asthma (ARIA) Guidelines - 2016 Revision. *J Allergy Clin Immunol.* 2017;140(4):950-8.
 27. Wallace DV, Dykewicz MS. Seasonal Allergic Rhinitis: A focused systematic review and practice parameter update. *Curr Opin Allergy Clin Immunol.* 2017;17(4):286-94.

- Accepted Article
28. Bosnic-Anticevich S, Kritikos V, Carter V, Yan KY, Armour C, Ryan D, et al. Lack of asthma and rhinitis control in general practitioner-managed patients prescribed fixed-dose combination therapy in Australia. *J Asthma*. 2018;55(6):684-94.
 29. Meltzer EO, Farrar JR, Sennett C. Findings from an Online Survey Assessing the Burden and Management of Seasonal Allergic Rhinoconjunctivitis in US Patients. *J Allergy Clin Immunol Pract*. 2017;5(3):779-89 e6.
 30. Collins JC, Schneider CR, Wilson F, de Almeida Neto AC, Moles RJ. Community pharmacy modifications to non-prescription medication requests: A simulated patient study. *Res Social Adm Pharm*. 2018;14(5):427-33.
 31. Kuehl BL, Abdulnour S, O'Dell M, Kyle TK. Understanding the role of the healthcare professional in patient self-management of allergic rhinitis. *SAGE Open Med*. 2015;3:2050312115595822.
 32. Prepageran N, Wang de Y, Nair G, Maurer M. The status quo and unmet needs in the management of allergic rhinitis and chronic rhinosinusitis: a Malaysian perspective. *Asia Pac Allergy*. 2014;4(3):142-8.
 33. Cote L, Normandeau M, Maheux B, Authier L, Lefort L. Collaboration between family physicians and community pharmacists: opinions of graduates in family medicine. *Can Fam Physician*. 2013;59(9):e413-20.
 34. Porteous T, Wyke S, Smith S, Bond C, Francis J, Lee AJ, et al. 'Help for Hay Fever', a goal-focused intervention for people with intermittent allergic rhinitis, delivered in Scottish community pharmacies: study protocol for a pilot cluster randomized controlled trial. *Trials*. 2013;14:217.
 35. Walls RS, Heddle RJ, Tang ML, Basger BJ, Solley GO, Yeo GT. Optimising the management of allergic rhinitis: an Australian perspective. *Med J Aust*. 2005;182(1):28-33.
 36. Smith L, Brown L, Saini B, Seeto C. Strategies for the management of intermittent allergic rhinitis: an Australian study. *Health Expect*. 2014;17(2):154-63.
 37. O'Connor J, Seeto C, Saini B, Bosnic-Anticevich S, Krass I, Armour C, et al. Healthcare professional versus patient goal setting in intermittent allergic rhinitis. *Patient Educ Couns*. 2008;70(1):111-7.
 38. Smith L, Nguyen T, Seeto C, Saini B, Brown L. The role of non-clinicians in a goal setting model for the management of allergic rhinitis in community pharmacy settings. *Patient Educ Couns*. 2011;85(2):e26-32.
 39. Lourenco O, Calado S, Sa-Sousa A, Fonseca J. Evaluation of allergic rhinitis and asthma control in a Portuguese community pharmacy setting. *J Manag Care Spec Pharm*. 2014;20(5):513-22.
 40. Lombardi C, Musicco E, Rastrelli F, Bettoncelli G, Passalacqua G, Canonica GW. The patient with rhinitis in the pharmacy. A cross-sectional study in real life. *Asthma Res Pract*. 2015;1:4.
 41. Kuipers E, Wensing M, de Smet P, Teichert M. Self-management research of asthma and good drug use (SMARAGD study): a pilot trial. *Int J Clin Pharm*. 2017;39(4):888-96.
 42. Canonica GW, Triggiani M, Senna G. 360 degree perspective on allergic rhinitis management in Italy: a survey of GPs, pharmacists and patients. *Clin Mol Allergy*. 2015;13:25.

43. Smith P, Price D, Harvey R, Carney AS, Kritikos V, Bosnic-Anticevich SZ, et al. Medication-related costs of rhinitis in Australia: a NostraData cross-sectional study of pharmacy purchases. *J Asthma Allergy*. 2017;10:153-61.
44. Bousquet J, Van Cauwenberge P, Khaltaev N. Allergic rhinitis and its impact on asthma. *J Allergy Clin Immunol*. 2001;108(5 Suppl):S147-334.
45. Agache I, Deleanu D, Khaltaev N, Bousquet J. [Allergic rhinitis and its impact upon asthma--update (ARIA 2008). Romanian perspective]. *Pneumologia*. 2009;58(4):255-8.
46. Bachert C, Jorissen M, Bertrand B, Khaltaev N, Bousquet J. Allergic Rhinitis and its impact on asthma update (ARIA 2008). The Belgian perspective. *B-ENT*. 2008;4(4):253-7.
47. Bousquet J, Farrell J, Crooks G, Hellings P, Bel EH, Bewick M, et al. Scaling up strategies of the chronic respiratory disease programme of the European Innovation Partnership on Active and Healthy Ageing (Action Plan B3: Area 5). *Clin Transl Allergy*. 2016;6:29.
48. Bousquet J, Schunemann HJ, Samolinski B, Demoly P, Baena-Cagnani CE, Bachert C, et al. Allergic Rhinitis and its Impact on Asthma (ARIA): achievements in 10 years and future needs. *J Allergy Clin Immunol*. 2012;130(5):1049-62.
49. Cagnani CE, Sole D, Diaz SN, Zernotti ME, Sisul JC, Borges MS, et al. [Allergic rhinitis update and its impact on asthma (ARIA 2008). Latin American perspective]. *Rev Alerg Mex*. 2009;56(2):56-63.
50. Kalayci O, Yorgancioglu A, Kalyoncu F, Khaltaev AN, Bousquet J. Allergic rhinitis and its impact on asthma update (ARIA 2008): the Turkish perspective. *Turk J Pediatr*. 2008;50(4):307-12.
51. Mullol J, Valero A, Alobid I, Bartra J, Navarro AM, Chivato T, et al. Allergic Rhinitis and its Impact on Asthma update (ARIA 2008). The perspective from Spain. *J Investig Allergol Clin Immunol*. 2008;18(5):327-34.
52. Pali-Scholl I, Pohl W, Aberer W, Wantke F, Horak F, Jensen-Jarolim E, et al. [Allergic rhinitis and its impact on asthma (ARIA update 2008) The Austrian perspective.]. *Wien Med Wochenschr*. 2009;159(3-4):87-92.
53. Pawankar R, Bunnag C, Chen Y, Fukuda T, Kim YY, Le LT, et al. Allergic rhinitis and its impact on asthma update (ARIA 2008)--western and Asian-Pacific perspective. *Asian Pac J Allergy Immunol*. 2009;27(4):237-43.
54. Yorgancioglu A, Kalayci O, Kalyoncu AF, Khaltaev N, Bousquet J. [Allergic rhinitis and its impact on asthma update (ARIA 2008). The Turkish perspective]. *Tuberk Toraks*. 2008;56(2):224-31.
55. Bousquet J, Hellings PW, Agache I, Bedbrook A, Bachert C, Bergmann KC, et al. ARIA 2016: Care pathways implementing emerging technologies for predictive medicine in rhinitis and asthma across the life cycle. *Clin Transl Allergy*. 2016;6:47.
56. Kotter J. *Leading change*. Boston, USA: Harvard Business School Press; 1996.
57. Bousquet J, Michel J, Standberg T, Crooks G, Iakovidis I, Gomez M. The European Innovation Partnership on Active and Healthy Ageing: the European Geriatric Medicine introduces the EIP on AHA Column. *Eur Geriatr Med*. 2014;5(6):361-2.

- Accepted Article
58. Bousquet J, Barbara C, Bateman E, Bel E, Bewick M, Chavannes NH, et al. AIRWAYS-ICPs (European Innovation Partnership on Active and Healthy Ageing) from concept to implementation. *Eur Respir J*. 2016;47(4):1028-33.
 59. Bousquet J, Neukirch F, Bousquet PJ, Gehano P, Klossek JM, Le Gal M, et al. Severity and impairment of allergic rhinitis in patients consulting in primary care. *J Allergy Clin Immunol*. 2006;117(1):158-62.
 60. Bousquet J, Annesi-Maesano I, Carat F, Leger D, Rugina M, Pribil C, et al. Characteristics of intermittent and persistent allergic rhinitis: DREAMS study group. *Clin Exp Allergy*. 2005;35(6):728-32.
 61. Bousquet PJ, Devillier P, Tadmouri A, Mesbah K, Demoly P, Bousquet J. Clinical Relevance of Cluster Analysis in Phenotyping Allergic Rhinitis in a Real-Life Study. *Int Arch Allergy Immunol*. 2015;166(3):231-40.
 62. del Cuvillo A, Montoro J, Bartra J, Valero A, Ferrer M, Jauregui I, et al. Validation of ARIA duration and severity classifications in Spanish allergic rhinitis patients - The ADRIAL cohort study. *Rhinology*. 2010;48(2):201-5.
 63. Jauregui I, Davila I, Sastre J, Bartra J, del Cuvillo A, Ferrer M, et al. Validation of ARIA (Allergic Rhinitis and its Impact on Asthma) classification in a pediatric population: the PEDRIAL study. *Pediatr Allergy Immunol*. 2011;22(4):388-92.
 64. Ozdalga E, Ozdalga A, Ahuja N. The smartphone in medicine: a review of current and potential use among physicians and students. *J Med Internet Res*. 2012;14(5):e128.
 65. Bourret R, Bousquet J, J M, T C, Bedbrook A, P D, et al. MASK rhinitis, a single tool for integrated care pathways in allergic rhinitis. *World Hosp Health Serv*. 2015;51(3):36-9.
 66. Bousquet J, Addis A, Adcock I, Agache I, Agusti A, Alonso A, et al. Integrated care pathways for airway diseases (AIRWAYS-ICPs). *Eur Respir J*. 2014;44(2):304-23.
 67. Bousquet J, Caimmi D, Bedbrook A, M Bewick, Hellings P, Devillier P, et al. Pilot study of mobile phone technology in allergic rhinitis in European countries. The MASK-rhinitis study *Allergy*. 2017:in press.
 68. Hellings PW, Muraro A, Fokkens W, Mullol J, Bachert C, Canonica GW, et al. A common language to assess allergic rhinitis control: results from a survey conducted during EAACI 2013 Congress. *Clin Transl Allergy*. 2015;5:36.
 69. Klimek L, Bergmann K, Biederman T, Bousquet J, Hellings P, al e. Visual analogue scales (VAS): measuring instruments for the documentation of symptoms and therapy monitoring in allergic rhinitis in everyday health care. Position Paper of the German Society of Allergology. *Allergo J Int*. 2017;26(1):16-24.
 70. Konig HH, Bernert S, Angermeyer MC, Matschinger H, Martinez M, Vilagut G, et al. Comparison of population health status in six european countries: results of a representative survey using the EQ-5D questionnaire. *Med Care*. 2009;47(2):255-61.

- Accepted Article
71. Smith AF, Pitt AD, Rodruiguez AE, Alio JL, Marti N, Teus M, et al. The economic and quality of life impact of seasonal allergic conjunctivitis in a Spanish setting. *Ophthalmic Epidemiol.* 2005;12(4):233-42.
 72. Bousquet J, Arnavielhe S, Bedbrook A, Fonseca J, Morais Almeida M, Todo Bom A, et al. The ARIA score of allergic rhinitis using mobile technology correlates with quality-of-life: The MASK study. *Allergy.* 2017;73(2):505-510.
 73. Azevedo P, Correia de Sousa J, Bousquet J, Bugalho-Almeida A, Del Giacco SR, Demoly P, et al. Control of Allergic Rhinitis and Asthma Test (CARAT): dissemination and applications in primary care. *Prim Care Respir J.* 2013;22(1):112-6.
 74. Fonseca JA, Nogueira-Silva L, Morais-Almeida M, Azevedo L, Sa-Sousa A, Branco-Ferreira M, et al. Validation of a questionnaire (CARAT10) to assess rhinitis and asthma in patients with asthma. *Allergy.* 2010;65(8):1042-8.
 75. Nogueira-Silva L, Martins SV, Cruz-Correia R, Azevedo LF, Morais-Almeida M, Bugalho-Almeida A, et al. Control of allergic rhinitis and asthma test--a formal approach to the development of a measuring tool. *Respir Res.* 2009;10:52.
 76. van der Leeuw S, van der Molen T, Dekhuijzen PN, Fonseca JA, van Gemert FA, Gerth van Wijk R, et al. The minimal clinically important difference of the control of allergic rhinitis and asthma test (CARAT): cross-cultural validation and relation with pollen counts. *NPJ Prim Care Respir Med.* 2015;25:14107.
 77. Bertsche T, Nachbar M, Fiederling J, Schmitt SP, Kaltschmidt J, Seidling HM, et al. Assessment of a computerised decision support system for allergic rhino-conjunctivitis counselling in German pharmacy. *Int J Clin Pharm.* 2012;34(1):17-22.
 78. Bousquet J, Reid J, van Weel C, Baena Cagnani C, Canonica GW, Demoly P, et al. Allergic rhinitis management pocket reference 2008. *Allergy.* 2008;63(8):990-6.
 79. Galimberti M, Passalacqua G, Incorvaia C, Castella V, Costantino MT, Cucchi B, et al. Catching allergy by a simple questionnaire. *World Allergy Organ J.* 2015;8(1):16.
 80. Guilemany JM, Garcia-Pinero A, Alobid I, Cardelus S, Centellas S, Bartra J, et al. Persistent allergic rhinitis has a moderate impact on the sense of smell, depending on both nasal congestion and inflammation. *Laryngoscope.* 2009;119(2):233-8.
 81. Bousquet J, Bewick M, Arnavielhe S, Mathieu-Dupas E, Murray R, Bedbrook A, et al. Work productivity in rhinitis using cell phones: The MASK pilot study. *Allergy.* 2017;72(10):1475-84.
 82. Bousquet J, Caimmi DP, Bedbrook A, Bewick M, Hellings PW, Devillier P, et al. Pilot study of mobile phone technology in allergic rhinitis in European countries: the MASK-rhinitis study. *Allergy.* 2017;72(6):857-65.
 83. Bousquet J, VandenPlas O, Bewick M, Arnavielhe S, Bedbrook A, Murray R, et al. Work Productivity and Activity Impairment Allergic Specific (WPAI-AS) Questionnaire using Mobile Technology: The MASK study. *J Investig Allergol Clin Immunol.* 2017:0.

- Accepted Article
84. Caimmi D, Baiz N, Tanno LK, Demoly P, Arnavielhe S, Murray R, et al. Validation of the MASK-rhinitis visual analogue scale on smartphone screens to assess allergic rhinitis control. *Clin Exp Allergy*. 2017;47(12):1526-1533.
 85. Cruz AA, Popov T, Pawankar R, Annesi-Maesano I, Fokkens W, Kemp J, et al. Common characteristics of upper and lower airways in rhinitis and asthma: ARIA update, in collaboration with GA(2)LEN. *Allergy*. 2007;62 Suppl 84:1-41.
 86. Bousquet J, Anto J, Demoly P, Schunemann H, Togias A, Akdis M, et al. Severe chronic allergic (and related) diseases: a uniform approach – a MeDALL-GA2LEN-ARIA Position Paper. *Int Arch Allergy Immunol*. 2012;158(3):216-31.
 87. Bousquet PJ, Combescure C, Neukirch F, Klossek JM, Mechin H, Daures JP, et al. Visual analog scales can assess the severity of rhinitis graded according to ARIA guidelines. *Allergy*. 2007;62(4):367-72.
 88. Schatz M, Meltzer EO, Nathan R, Derebery MJ, Mintz M, Stanford RH, et al. Psychometric validation of the rhinitis control assessment test: a brief patient-completed instrument for evaluating rhinitis symptom control. *Ann Allergy Asthma Immunol*. 2010;104(2):118-24.
 89. Demoly P, Jankowski R, Chassany O, Bessah Y, Allaert FA. Validation of a self-questionnaire for assessing the control of allergic rhinitis. *Clin Exp Allergy*. 2011;41(6):860-8.
 90. Morais-Almeida M, Pite H, Pereira AM, Todo-Bom A, Nunes C, Bousquet J, et al. Prevalence and classification of rhinitis in the elderly: a nationwide survey in Portugal. *Allergy*. 2013;68(9):1150-7.
 91. Klimek L, Bachert C, Mosges R, Munzel U, Price D, Virchow JC, et al. Effectiveness of MP29-02 for the treatment of allergic rhinitis in real-life: results from a noninterventional study. *Allergy Asthma Proc*. 2015;36(1):40-7.
 92. Bousquet J, Bachert C, Canonica GW, Mullol J, Van Cauwenberge P, Bindeslev Jensen C, et al. Efficacy of desloratadine in intermittent allergic rhinitis: a GALEN study. *Allergy*. 2009;64(15):16-23.
 93. Bousquet J, Bachert C, Canonica GW, Mullol J, Van Cauwenberge P, Jensen CB, et al. Efficacy of desloratadine in persistent allergic rhinitis - a GA(2)LEN study. *Int Arch Allergy Immunol*. 2010;153(4):395-402.
 94. Klimek L, Bergmann KC, Biedermann T, Bousquet J, Hellings P, Jung K, et al. Visual analogue scales (VAS): Measuring instruments for the documentation of symptoms and therapy monitoring in cases of allergic rhinitis in everyday health care: Position Paper of the German Society of Allergology (AeDA) and the German Society of Allergy and Clinical Immunology (DGAKI), ENT Section, in collaboration with the working group on Clinical Immunology, Allergology and Environmental Medicine of the German Society of Otorhinolaryngology, Head and Neck Surgery (DGHNOKHC). *Allergo J Int*. 2017;26(1):16-24.
 95. Scadding GK, Kariyawasam HH, Scadding G, Mirakian R, Buckley RJ, Dixon T, et al. BSACI guideline for the diagnosis and management of allergic and non-allergic rhinitis (Revised Edition 2017; First edition 2007). *Clinical and Experimental Allergy*. 2017;47:856-89.
 96. Compalati E, Braidò F, Canonica GW. Sublingual Immunotherapy: Recent Advances. *Allergology International*. 2013;62(4):415-23.

97. Passalacqua G, Durham SR, Global A, Asthma European N. Allergic rhinitis and its impact on asthma update: allergen immunotherapy. *J Allergy Clin Immunol*. 2007;119(4):881-91.
98. Passalacqua G, Garelli V, Canonica GW. Sublingual immunotherapy for allergic rhinitis and conjunctivitis. *Immunotherapy*. 2013;5(3):257-64.
99. Ozturk AB, Celebioglu E, Karakaya G, Kalyoncu AF. Protective efficacy of sunglasses on the conjunctival symptoms of seasonal rhinitis. *Int Forum Allergy Rhinol*. 2013;3(12):1001-6.
100. O'Meara TJ, Sercombe JK, Morgan G, Reddel HK, Xuan W, Tovey ER. The reduction of rhinitis symptoms by nasal filters during natural exposure to ragweed and grass pollen. *Allergy*. 2005;60(4):529-32.
101. Raulf M, Buters J, Chapman M, Cecchi L, de Blay F, Doekes G, et al. Monitoring of occupational and environmental aeroallergens-- EAACI Position Paper. Concerted action of the EAACI IG Occupational Allergy and Aerobiology & Air Pollution. *Allergy*. 2014;69(10):1280-99.
102. Garavento W, Di Berardino F, Romagnoli M, Sambataro G, Gaini RM. Nasal rinsing with hypertonic solution: an adjunctive treatment for pediatric seasonal allergic rhinoconjunctivitis. *Int Arch Allergy Immunol*. 2005;137(4):310-4.
103. Jeffe JS, Bhushan B, Schroeder JW, Jr. Nasal saline irrigation in children: a study of compliance and tolerance. *Int J Pediatr Otorhinolaryngol*. 2012;76(3):409-13.
104. Bender BG. Motivating patient adherence to allergic rhinitis treatments. *Curr Allergy Asthma Rep*. 2015;15(3):10.
105. Tay SY, Chao SS, Mark KT, Wang Y. Comparison of the distribution of intranasal steroid spray using different application techniques. *Int Forum Allergy Rhinol*. 2016;6(11):1204-10.
106. Ganesh V, Banigo A, McMurran AEL, Shakeel M, Ram B. Does intranasal steroid spray technique affect side effects and compliance? Results of a patient survey. *J Laryngol Otol*. 2017;131(11):991-6.
107. Bousquet J, Onorato GL, Bachert C, Barbolini M, Bedbrook A, Bjermer L, et al. CHRODIS criteria applied to the MASK (MACVIA-ARIA Sentinel Network) Good Practice in allergic rhinitis: a SUNFRAIL report. *Clin Transl Allergy*. 2017;7:37.
108. Caimmi D, Baiz N, Tanno LK, Demoly P, Arnavielhe S, Murray R, et al. Validation of the MASK-rhinitis visual analogue scale on smartphone screens to assess allergic rhinitis control. *Clin Exp Allergy*. 2017;47(12):1526-33.
109. Bousquet J, Agache I, Aliberti MR, Angles R, Annesi-Maesano I, Anto JM, et al. Transfer of innovation on allergic rhinitis and asthma multimorbidity in the elderly (MACVIA-ARIA) - EIP on AHA Twinning Reference Site (GARD research demonstration project). *Allergy*. 2018;73(1):77-92.
110. Bousquet J, Arnavielhe S, Bedbrook A, Alexis-Alexandre G, Eerd Mv, Murray R, et al. Treatment of allergic rhinitis using mobile technology with real world data: The MASK observational pilot study. *Allergy*. 2018;73(9):1763-1774.

- Accepted Article
111. Bousquet J, Arnavielhe S, Bedbrook A, Fonseca J, Morais Almeida M, Todo Bom A, et al. The Allergic Rhinitis and its Impact on Asthma (ARIA) score of allergic rhinitis using mobile technology correlates with quality of life: The MASK study. *Allergy*. 2018;73(2):505-10.
 112. Treherne JM, Young JM. [3H]-(+)-N-methyl-4-methyldiphenhydramine, a quaternary radioligand for the histamine H1-receptor. *Br J Pharmacol*. 1988;94(3):797-810.